

Central serous chorio- retinopathy (CSCR)

Central serous chorio-retinopathy (CSCR) is a condition that affects the retina- the light sensitive tissue that lines the back of the eye.




Diagram of the eye

Central serous chorio-retinopathy (CSCR), refers to a collection of fluid under the retina. This is caused by a disturbance in the pumping action of special cells called RPE cells (retinal pigment epithelial cells) and/or abnormalities in the vascular (blood vessel) layer, known as choroid.

RPE cell layer and blood vessel layer (choroid) line the outer surface of retina and both layers function to keep the

retina healthy. This dysfunction results in fluid leakage under the retina in a bubble-like swelling called central serous chorio-retinopathy (CSCR).

What are the symptoms of CSCR?

CSCR usually affects those between 30 to 50 years of age. Males are more likely to be affected than females. Main symptoms include:

- painless blurriness of central vision.
- distortion or a change in size of an object.
- straight objects or lines seeming curved (distorted).
- difficulty in reading small prints with the affected eye.
- some patients notice that they need to change reading glasses more often.

In most cases, the symptoms only last for a few months; however it may last longer in a minority of patients, potentially leading to a long-term worsening of vision.

What causes CSCR?

The exact factors that cause the development of CSCR related fluid

have not yet been determined. However, we know of various risk factors that are associated with the development of CSCR.

1) Steroids

Steroids are a known risk factor for the development of CSCR. These include steroids in different forms such as inhalers for asthma, nasal spray for hay fever, steroid cream for eczema and steroid tablets such as prednisolone. Rarely, in a condition (known as Cushing syndrome), an over production of the body's natural steroid hormone could lead to the development of CSCR.

2) Stress

A major stressful event, either work-related or personal, is believed to trigger the development of CSCR in some patients.

3) Psychological make-up

Research has shown that certain personality types (particularly those who are hard-driven and competitive), are more at risk of developing this condition.

4) Genetic risk

There are ongoing studies suggesting some patients may have changes in certain genes that can trigger CSCR when exposed to certain environmental factors.

How is CSCR diagnosed?


Generally, the diagnosis is made by taking your detailed history which includes medical and drug history,

clinical examination of eyes and imaging tests.


Types of imaging tests:

1) Optical coherence tomography

(OCT) is a scan of the retina. It is a non-invasive camera-based imaging test which uses light waves to take cross-section pictures of your retina; It is used to identify the fluid under the retina, along with detailed structural changes secondary to CSCR. This imaging tool is not only useful in making a diagnosis but is also helpful in monitoring your condition. Some forms of OCT (OCT- angiography) look at abnormal blood vessel growth that can rarely complicate CSCR. Your consulting eye doctor will decide whether this test is needed.



Normal retina on OCT imaging



OCT image showing retinal fluid in CSCR

2) FFA and ICG test (Fundus fluorescein angiography (FFA) and Indocyanine green angiography (ICG))



risk of vision loss) and not everyone is suitable for the treatment.

2. Conventional laser:

If the source of the retinal fluid associated with CSCR is away from the central macular then this treatment option may be considered.

3. Micro-pulse laser

This is a sub-threshold (low -power) laser which is applied in short pulses to the area of leakage, resulting in fixing the retinal fluid. Recent studies have shown that this treatment is inferior to PDT.

4. Oral medications

There are some recent studies that have shown the effectiveness of certain tablets in reducing the fluid in CSCR. Eplerenone and Spironolactone are mineralocorticoid inhibitors, commonly used to lower blood pressure or treat heart failure. These medications have shown an improvement in CSCR symptoms for some patients, although other studies have not shown benefit. Further studies are currently being carried out to evaluate the role of these treatments. When on this treatment, these drugs may cause side effects in some patients including changing the salt (potassium) level in the blood which will need monitoring. Other potential side effects which vary by drug will be discussed by your doctor during the clinic consultation.

How can I get treatment?

As mentioned previously, all the above treatments listed are not licensed or

NICE approved for managing CSCR. This means your eye doctor may need to try and get funding for treatment and the success of this depends on your local health authority funding budget. This situation can often lead to delays in treatment or in some cases it is not possible to get funding for a recommended treatment. The funding situation is currently subject to change and you will need to discuss this further with your eye doctor.

Can changing glasses help manage CSCR?

Changing existing glasses or getting new glasses does not help manage CSCR, as your glasses prescription can be different depending on the amount of retinal fluid you have. Changing your glasses prescription would only help temporarily; therefore, it is advisable to wait until the retinal fluid has been completely resolved.

Authors: Yasir Khan, Adam Mapani, Niaz Islam, Adnan Tufail

Revision number: 1

Approval date: September 2018

Review date: September 2020

Moorfields Eye Hospital NHS Foundation Trust
City Road, London EC1V 2PD
Phone: 020 7253 3411
www.moorfields.nhs.uk

Moorfields Direct telephone helpline
Phone: 020 7566 2345
Monday-Friday, 8.30am-9pm
Saturday, 9am-5pm





Information and advice on eye conditions and treatments from experienced ophthalmic-trained nurses.

Patient advice and liaison service (PALS)

Phone: 020 7566 2324/ 020 7566 2325
Email: moorfields.pals@nhs.net
Moorfields' PALS team provides confidential advice and support to help you with any concerns you may have about the care we provide, guiding you through the different services available at Moorfields. The PALS team can also advise you on how to make a complaint.

Your right to treatment within 18 weeks

Under the NHS constitution, all patients have the right to begin consultant-led treatment within 18 weeks of being referred by their GP. Moorfields is committed to fulfilling this right, but if you feel that we have failed to do so, please contact our patient advice and liaison service (PALS) who will be able to advise you further (see above). For more information about your rights under the NHS constitution, visit www.nhs.uk/choiceinthenhs